

Erratum: Elements of a theory for multiparton interactions in QCD

Markus Diehl,^a Daniel Ostermeier^b and Andreas Schäfer^b

^aDeutsches Elektronen-Synchrotron DESY, 22603 Hamburg, Germany

^bInstitut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany

E-mail: markus.diehl@desy.de,

Daniel.Ostermeier@physik.uni-regensburg.de,

Andreas.Schaefer@physik.uni-regensburg.de

ERRATUM TO: [JHEP03\(2012\)089](#)

ARXIV EPRINT: [1111.0910](#)

1. In equations (3.41) and (3.42), all terms v^+v^- should be changed into v^+w^- .

2. The second part of equation (4.25) correctly reads

$$M_{a_1, a_2}^{n_1, n_2}(y^2) = \frac{1}{2}(p^+)^{1-n_1-n_2} \int dy^- \langle p | \mathcal{O}_{a_1}^{+\cdots+}(0) \mathcal{O}_{a_2}^{+\cdots+}(y) | p \rangle_{y^+=0}$$

with a factor 1/2 instead of 2.

3. Equation (5.75) correctly reads

$$\left[\prod_{i=1}^2 \int d^2 \mathbf{k}_i \right] {}^1 F_{a_1, \bar{a}_2}(x_i, \mathbf{k}_i, \mathbf{r}) \Big|_{g \rightarrow q\bar{q}} = \frac{\alpha_s}{4\pi^2} \frac{1}{x_1 + x_2} f_1^g(x_1 + x_2) T_{a_1, \bar{a}_2}^{ll'} \left(\frac{x_1}{x_1 + x_2} \right) \times \int d^2 \mathbf{k} \frac{(\mathbf{k} + \frac{1}{2}\mathbf{r})^l (\mathbf{k} - \frac{1}{2}\mathbf{r})^{l'}}{(\mathbf{k} + \frac{1}{2}\mathbf{r})^2 (\mathbf{k} - \frac{1}{2}\mathbf{r})^2},$$

i.e. the factors $(2\pi)^2$ should be omitted on both sides.

4. Together with the surrounding text, equation (5.78) for the double parton distribution in $4 - 2\epsilon$ dimensions should read:

The result for $F_{q, \bar{q}}(x_i, \mathbf{k}_i, \mathbf{r})$ is then the same as in (5.42) with a modified kernel

$$T_{q, \bar{q}}^{ll'}(u; \epsilon) = \delta^{ll'} [u^2 + (1-u)^2 - \epsilon]/(1-\epsilon)$$

times a power of $(2\pi)^\epsilon$ we need not specify here.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.