

RESEARCH ARTICLE

Open Access



Synthesis, characterization, molecular docking, analgesic, antiplatelet and anticoagulant effects of dibenzylidene ketone derivatives

Tauqeer Ahmed¹, Arif-ullah Khan^{1*}, Muzaffar Abbass^{1,5}, Edson Rodrigues Filho², Zia Ud Din^{2,3} and Aslam Khan⁴

Abstract

In this study dibenzylidene ketone derivatives (2*E*,5*E*)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a) and (1*E*,4*E*)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl) oct-1-en-3-one (AK-2a) were newly synthesized, inspired from curcuminoids natural origin. Novel scheme was used for synthesis of AK-1a and AK-2a. The synthesized compounds were characterized by spectroscopic techniques. AK-1a and AK-2a showed high computational affinities (E-value > -9.0 kcal/mol) against cyclooxygenase-1, cyclooxygenase-2, proteinase-activated receptor 1 and vitamin K epoxide reductase. AK-1a and AK-2a showed moderate docking affinities (E-value > -8.0 kcal/mol) against mu receptor, kappa receptor, delta receptor, human capsaicin receptor, glycoprotein IIb/IIIa, prostacyclin receptor I₂, antithrombin-III, factor-II and factor-X. AK-1a and AK-2a showed lower affinities (E-value > -7.0 kcal/mol) against purinoceptor-3, glycoprotein-VI and purinergic receptor P₂Y₁₂. In analgesic activity, AK-1a and AK-2a decreased numbers of acetic acid-induced writhes ($P < 0.001$ vs. saline group) in mice. AK-1a and AK-2a significantly prolonged the latency time of mice ($P < 0.05$, $P < 0.01$ and $P < 0.001$ vs. saline group) in hotplate assay. AK-1a and AK-2a inhibited arachidonic acid and adenosine diphosphate induced platelet aggregation with IC₅₀ values of 65.2, 37.7, 750.4 and 422 μM respectively. At 30, 100, 300 and 1000 μM concentrations, AK-1a and AK-2a increased plasma recalcification time ($P < 0.001$ and $P < 0.001$ vs. saline group) respectively. At 100, 300 and 1000 μg/kg doses, AK-1a and AK-2a effectively prolonged bleeding time ($P < 0.001$ and $P < 0.01$ vs. saline group) respectively. Thus in-silico, in-vitro and in-vivo investigation of AK-1a and AK-2a reports their analgesic, antiplatelet and anticoagulant actions.

Keywords: Dibenzylidene ketone derivatives, Computational studies, Analgesic, Antiplatelet, Anticoagulant, Arachidonic acid

Introduction

Pain is an unfavorable sensory and emotional experience that is associated with the potential tissue damage and explained in terms of such damage [1]. Noxious effects such as ulceration, gastrointestinal bleeding by non-steroidal anti-inflammatory drugs and drowsiness, nausea and tolerance by opiates usage limits their use in management of pain [2]. Platelets play vital role in a complex

processes which are involved in haemostasis and thrombosis [3]. The most common cause of peripheral artery diseases (PAD) is atherosclerosis and such patients have more chance of myocardial infarction, stroke or death with cardiovascular events and it is 3:1 in comparison to persons without PAD [4]. Antiplatelet agents are used in management of arterial thrombosis. Moreover, anticoagulants inhibit proteases in coagulation cascade [5]. Interference in natural balance among pro-coagulant and anticoagulant due to genetic or any other acquired factors may results in bleeding or thrombotic disorders. Thrombin is a key enzyme of coagulation cascade which has many significant biological functions including platelet

*Correspondence: arif.ullah@riphah.edu.pk

¹ Riphah Institute of Pharmaceutical Sciences, Riphah International University, Islamabad, Pakistan

Full list of author information is available at the end of the article



activation, fibrinogen conversion to fibrin network and feedback amplification of coagulation. Different tissue factors are involved in thrombus formation in order to prevent hemorrhage [6]. Coagulation cascade involves intrinsic and extrinsic pathways [7]. The former has a role in the growth and maintenance of fibrin while the latter plays its role in the initiation of fibrin formation. Extrinsic pathway requires tissue factors for its activation which after vascular injury becomes exposed to the blood which ultimately results in thrombin activation [8]. Among antiplatelet agent and anticoagulant drugs which are available commercially, for thrombotic disorders, these agents are associated with certain limitations and side effects [9]. Chemically curcumin is 1,7-bis (4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione. It is a yellow-orange colored pigment which is derived from the rhizome of *Curcuma longa* [10]. The plant has a wide spectrum of pharmacological properties and traditionally it has been used for many ailments since centuries [11]. The reported activities of curcumin are antioxidant, anti-inflammatory, antitumor, antibacterial, antifungal and antiviral [10]. Curcumin also showed inhibition in platelet aggregation and antithrombotic effects [12, 13]. Concerning structural aspects, dibenzylidene ketone moieties are considered curcumin analogues, which are compounds of great importance. Structurally, curcuminoids contain two aryl rings connected at the ends of a C₇ carbon-chain where a dienone composes an extended conjugated system. Dibenzylidene ketone derivatives also contain a dienone system connecting two aryl groups at the ends of a C₅ carbon chain. Dienones are good Michael acceptors, allowing its reaction with important biomolecules interfering in biological processes. Previous reported activities of dibenzylidene ketone derivatives include antiparasitic activity, cytotoxicity, antimicrobial activity, analgesic activity [14–16]. Based on previous literature studies, two novel dibenzylidene ketone derivatives i.e. (2*E*,5*E*)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a) and (1*E*,4*E*)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl) oct-1-en-3-one (AK-2a) were synthesized and characterized. AK-1a and AK-2a were investigated for their analgesic, antiplatelet and anticoagulant effects using different pharmacological and computational assays.

Materials and methods

Chemicals

Adenosine diphosphate (ADP) and arachidonic acid (AA) were purchased from Chrono-Log association. Benzaldehyde, cyclopentanone, dimethyl sulfoxide, ethanol and methoxybenzaldehyde were purchased from Merck Millipore., Billerica, MA, USA. Aspirin, calcium chloride (CaCl₂), diclofenac sodium, heparin, phosphate buffers

solution (PBS) and sodium citrate were obtained from Sigma chemicals., Dt. Louis, MO, USA. The tramadol was acquired from Searle Karachi-Pakistan. All chemicals used were of analytical grade.

Animals

Balb-C mice (25–30 g) of both gender were utilized for this study. All animals were housed according to the standard protocols 25 ± 2 °C, 12 h duration of natural light and dark cycle. Healthy diet was given to mice and water ad libitum. The study was performed in accordance with protocols of Institute of Laboratory Animal Resources, Commission on Life Sciences University, National Research Council (1996) and approved by Riphah Institute of Pharmaceutical Sciences (RIPS) Ethical Committee (Reference No: REC/RIPS/2016/009).


Synthesis of AK-1a and AK-2a

Novel way of synthesis was carried out. The monoarylidene derivative was synthesized by the reaction of cyclopentanone with *p*-methoxy benzaldehyde. DIMCARB was utilized as a catalyst in this reaction. DIMCARB was used in catalytic amount to obtain selective monoarylidene cyclic derivative in a green solvent (EtOH:H₂O), further second step leads to get an unsymmetrical bis-(arylmethylidene)-cycloalkanones. The synthesis of compound was carried out at room temperature from the reaction of intermediate 1 with *p*-nitro benzaldehyde. The scheme of the synthesized compound along with its structure is shown in Fig. 1. Chemical characterization was carried out based on the analysis of spectroscopic data. Fourier transform mass spectrometry (FTMS) of AK-1a as shown in Fig. 2. Synthesis of AK-2a was carried in a two-step reaction. In the first step, cycloalkanone was reacted with an aldehyde in a DIMCARB catalysed reaction, while in the second step monoarylidene derivative was reacted with the aldehyde through Knoevenagel condensation to get the required product. DIMCARB can be recovered by distillative dissociation–reassociation process in a vacuum or under an atmosphere of CO₂. The 2-heptanone was reacted with *p*-nitro benzaldehyde in an acidic medium to get intermediate, and then intermediate yield AK-2a. The scheme of the novel synthesized compound AK-2a along with its structure is shown in Fig. 1. Chemical characterization was carried out based on the analysis of spectroscopic data. Fourier transform mass spectrometry (FTMS) of AK-2a as shown in Fig. 3.

Spectral analysis

AK-1a

Percent yield: 84. Decompose at: 96–98 °C. ¹H NMR (400 MHz, CDCl₃) δ 7.57 (d, J = 8.8 Hz, 3H), 7.52 (d,



$J = 8.4$ Hz, 3H), 7.40 (d, $J = 8.6$ Hz, 2H), 6.97 (d, $J = 8.9$ Hz, 2H), 3.86 (s, 3H), 3.08 (s, 4H). ^{13}C NMR (101 MHz, CDCl_3) δ 196.21 (1C), 160.90 (1C), 138.27 (1C), 135.28 (1C), 134.71 (1C), 134.32 (1C), 132.73 (2C), 131.93 (2C), 129.16 (2C), 128.70 (1C), 114.51 (2C), 77.48 (1C), 76.84 (1C), 55.54 (1C), 26.60 (2C). HRMS ESI(+): calcd for $\text{C}_{20}\text{H}_{17}\text{NNaO}_4^+$ ($M + \text{Na}$) 358.1050, found 358.1044.


AK-2a

Percent yield: 80. m.p: 180.5–181.5 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.28 (dd, $J = 8.5, 7.7$ Hz, 4H), 7.80–7.70 (m, 3H), 7.56 (d, $J = 8.7$ Hz, 2H), 7.49 (s, 1H), 7.43 (d, $J = 15.7$ Hz, 1H), 2.65–2.58 (m, 2H), 1.53–1.42 (m, 2H), 1.37 (dd, $J = 14.7, 7.2$ Hz, 2H), 0.91 (t, $J = 7.2$ Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 191.74 (1C), 148.73 (1C), 147.58 (1C), 146.65 (1C), 142.33 (1C), 141.43 (1C), 141.07 (1C), 135.88 (1C), 129.97 (2C), 129.03 (2C), 125.92 (1C), 124.38 (2C), 124.01 (2C), 31.22 (1C), 27.24 (1C), 23.03 (1C), 13.94 (1C). HRMS ESI(+): calcd for $\text{C}_{21}\text{H}_{20}\text{N}_2\text{NaO}_5^+$ ($M + \text{Na}$) 403.1264, found 403.1240.

In-silico study

Molecular docking is an informative tool which is used to investigate the affinity between ligand and protein

targets. We used Auto Dock Vina program for docking study through PyRx [17, 18]. Affinity of best docked pose of ligand and protein target complex was determined by E-value (kcal/mol). It provides prediction of binding free energy and binding constant for docked ligands [19]. 3D-structures of test compounds (AK-1a and AK-2a) were prepared in discovery studio visualiser (DSV) and saved as PDB format. 3D-structures of target proteins were taken from <http://www.rcsb.org/pdb/home/home.do>. The target proteins involved in pain pathways are cyclooxygenase-1 (COX-1, PDB-ID: 3N8X), cyclooxygenase-2 (COX-2, PDB-ID: 1PXX), mu receptor (PDB-ID: 5C1M), kappa receptor (PDB-ID: 4DJH), delta receptor (PDB-ID: 4EJH), human capsaicin receptor (HCR, PDB-ID: 3J9J) and purinoceptor-3 (P2X3, PDB-ID: 5SVL). The target proteins involved in platelet aggregation are glycoprotein-IIb/IIIa (GP-IIb/IIIa, PDB-ID: 2VDM), glycoprotein-VI (GP-VI, PDB-ID: 2G17), purinergic receptor (P_2Y_{12} , PDB-ID: 4PXZ), prostacyclin receptor I_2 (PG- I_2 , PDB-ID: 4F8K) and proteinase-activated receptor 1 (PAR-1, PDB-ID: 3VW7). The target proteins involved in blood coagulation process are antithrombin-III (AT-III, PDB-ID: 2B4X), factor-II (F-II, PDB-ID: 1KSN), factor-IX (F-IX, PDB-ID: 1XMN),



factor-X (F-X, PDB-ID: 1RFN) and vitamin-K epoxide reductase (VKOR, PDB-ID: 3KP9). All the target proteins were then purified by removing ligands and other entities which might occupy nearby space using Biovia Discovery Studio Client 2016. The structures of standard drug molecules were downloaded from pubchem data base (<https://pubchem.ncbi.nlm.nih.gov/search/>). Reference analgesic drugs are aspirin (PubChem CID: 2244), morphine (PubChem CID: 5288826) and capsaizepine (PubChem CID: 2733484). Standard antiplatelet drugs are aspirin (PubChem CID: 2244), tirofiban (PubChem CID: 60947), hinokitiol (PubChem CID: 3611), clopidogrel (PubChem CID: 10066813), beraprost (PubChem CID: 6917951) and vorapaxar (PubChem CID: 10077130). Reference anti-coagulant drugs are heparin sulphate (PubChem CID: 53477714), apixaban (PubChem CID: 10182969), argatroban (PubChem CID: 92722), pegnivacogin (PubChem CID: 86278323) and warfarin (PubChem CID: 54678486). All these structures were downloaded in .xml format and converted to PDB format via Open Babel JUI software. PDB form of both ligand and standard as well as target proteins were converted to PDBQT via AutoDockTools (Version1.5.6 Sep_17_14) where add kollman charges and compute gasteiger charges were added and Ad4 type was


assigned. Both the test compounds along with protein targets in PDBQT form were loaded in software named as PyRx and then docked against the respective targets. Binding affinity was calculated and shown in kcal/mol. For post docking interaction Discovery studio visualizer was used for number of hydrogen bonds (classical and non-classical) and binding amino acid residues: alanine (ALA), asparagine (ASN), arginine (ARG), aspartic acid (ASP), cysteine (CYS), glutamine (GLN), glutamic acid (GLU), glycine (GLY), histidine (HIS), leucine (LEU), lysine (LYS), serine (SER), threonine (THR), tryptophan (TRP), tyrosine (TYR), valine (VAL) and phenylalanine (PHE) showed in the form 2D interaction.

Analgesic models

The analgesic activity was carried out by using two standard protocols i.e. acetic acid-induced writhing test and hot plate test in order to evaluate the peripheral and central effects of analgesia.

Acetic acid-induced writhing test

Mice were divided into five different groups, having five mice in each. After 30 min writhing were induced by an IP injection of 0.1 mL of 0.7% (by volume) acetic acid



solution [20]. Drug pretreatment times were chosen so that writhing was counted over a period of maximum analgesic activity. AK-1a and AK-2a in a dose-dependent manner (0.5–100 mg/kg) decreased acetic acid-induced writhes injected through intraperitoneal (IP) route. Perception of pain was recorded in the form of abdominal constrictions and stretches of the hind limb called as a writhe. Some mice showed half writhe. Two half writhes were considered as equal to one full writhe. The writhing episodes were recorded for 20 min. Control group was administered with normal saline (10 mL/kg). Diclofenac sodium was used as a positive control.

Hot plate test

The latency period of the test compounds were evaluated by hot plate assay according to the protocols as previously used with little modifications [21]. Mice were divided into five different groups, having five mice in each. The animals were placed individually on the

hot plate (55 ± 2 °C) and the observations (jumping or licking paws) were recorded at 30, 60, 90 and 120 min. Normal saline (10 mL/kg) was given to control group, tramadol (30 mg/kg) was used as a positive control.

Antiplatelet assay

Antiplatelet activity was performed to check whether the test compounds possess any effect on platelet aggregation. It was determined by whole-blood aggregometry method, which was performed by an impedance aggregometer (Model 591, Chrono-Log) as previously described [22]. Arterial or venous blood samples were collected from healthy volunteers in plastic tubes having 3.2% sodium citrate anticoagulant (9:1). Measurements were performed at 37 °C and 1200 rpm stirring speed. According to the manufacturer recommendations, 0.5 mL of citrated blood was diluted with same volume of normal saline (0.9%) which was prewarmed for 5 min at 37 °C. 30 μ L, AK-1a and AK-2a at 1, 3, 10, 30, 100, 300, 1000 μ M concentrations were also added to the tube. After placing the electrode, aggregation was induced by

different agonists like AA (1.5 mM) and ADP (10 μ M). Platelet aggregation response was continually monitored for 6 min as an electrical impedance in ohms. Then mean percent platelet inhibition was calculated. Aspirin was used as positive control.

Anticoagulant activity

Anticoagulant activity of the test compounds were performed using following experiments.

Plasma recalcification time (PRT)

Anticoagulant potential of the test compounds were determined by PRT method [23]. The blood samples were obtained from healthy volunteers in tubes containing 3.8% sodium citrate (9:1) in order to prevent the clotting process. Centrifugation (15 min at rate 3000 rpm) was carried out in order to obtain platelet poor plasma. 0.2 mL plasma, 0.1 mL of different concentration of the test compounds (30, 100, 300 and 1000 μ M) and 0.3 mL of CaCl_2 (25 mM) were then added together in a clean fusion tube and incubated in a water bath at 37 °C. Heparin (440 μ M) was used as positive control. The clotting time was recorded with a stopwatch by tilting the test tubes every 5 s.

Bleeding time (BT)

Anticoagulant potential of AK-1a and AK-2a was also determined by in-vivo tail BT method in mice [24]. AK-1a and AK-2a (100, 300 and 1000 μ g/kg) were administered intravenously via tail vein of mice. After 10 min mice were anesthetized using diethyl ether and a sharp cut (3 mm) deep at tip of the tail were made. The tail was then immersed into PBS which was pre warmed to 37 °C. BT was recorded from the time when bleeding started till it stopped completely observation was made up-to 10 min. Heparin (40 μ g/kg) was utilized as a positive control.

Statistical analysis

Data were expressed as mean \pm standard error of mean (SEM) and analyzed by using one-way analysis of variance (ANOVA), with post hoc Tukey's test. Data were considered significant at $P < 0.05$. Bar graphs were analyzed using Graph Pad Prism (GraphPad, San Diego, CA, USA).

Results

Molecular docking evaluation

The results of E-values, hydrogen bonds and binding residues of AK-1a and AK-2a with target proteins involved in pain pathways along with standard drugs are shown in Table 1 and Figs. 4, 5, 6, 7. The results of E-values, hydrogen bonds and binding residues of AK-1a and AK-2a with

target proteins involved in platelet aggregation along with standard drugs are shown in Table 2 and Figs. 7, 8, 9. The results of E-values, hydrogen bonds and binding residues of AK-1a and AK-2a with target proteins involved in coagulation process along with the standard drugs are shown in Table 3 and Figs. 10, 11, 12.

Effect on acetic acid-induced writhings

Saline group (10 mL/kg) showed 88 ± 2.28 numbers of writhes. The writhes count in AK-1a treated group (1, 10, 20, 30 and 100 mg/kg) were decreased to 77 ± 1.51 , 60.60 ± 1.07 , 51.80 ± 0.73 , 42.40 ± 1.72 and 33.80 ± 1.20 ($P < 0.001$ vs. saline group) respectively. Diclofenac sodium (20 mg/kg) decreased numbers of writhes to 29.80 ± 1.77 ($P < 0.001$ vs. saline group). AK-2a showed significant response in acetic acid induced writhing. The writhes count in AK-2a treated group (0.5, 1, 3 and 5 mg/kg) were decreased to 52.40 ± 1.40 , 38.60 ± 1.20 , 32.60 ± 1.50 and 2.00 ± 1.26 ($P < 0.001$ vs. saline group) respectively as shown in Fig. 13.



Effect on latency time

The latency time of saline group (10 mL/kg) at 0, 30, 60, 90 and 120 min were 7.35 ± 0.12 , 8.33 ± 0.13 , 8.56 ± 0.10 , 8.71 ± 0.10 and 8.70 ± 0.03 s respectively. AK-1a dose dependently (1, 10, 20 and 30 mg/kg) prolonged latency time against thermal pain generation. The latency time of AK-1a (1 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 5.42 ± 0.15 , 6.28 ± 0.15 , 8.63 ± 0.28 , 10.40 ± 0.19 , 11.47 ± 0.27 s ($P < 0.001$ vs. saline group) respectively. The latency time of AK-1a (10 mg/kg) treated group at 0, 30, 60, 90, 120 min were 5.78 ± 0.28 , 7.51 ± 0.20 , 9.36 ± 0.32 , 10.71 ± 0.39 and 12.80 ± 0.24 s ($P < 0.001$ vs. saline group) respectively. The latency time of AK-1a (20 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 7.33 ± 0.29 , 8.81 ± 0.26 , 9.27 ± 0.33 , 11.81 ± 0.24 and 16.92 ± 0.55 s ($P < 0.001$ vs. saline group) respectively. The latency time of AK-1a (30 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 9.69 ± 0.31 , 10.40 ± 0.36 , 11.88 ± 0.13 , 13.67 ± 0.23 and 15.66 ± 0.33 s ($P < 0.001$ vs. saline group) respectively. The latency time of tramadol (30 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 7.33 ± 0.20 , 13.07 ± 0.18 , 13.97 ± 0.12 , 14.69 ± 0.27 and 15.61 ± 0.18 s ($P < 0.001$ vs. saline group) respectively as shown in Fig. 14. AK-2a dose dependently (0.5, 1, 3 and 5 mg/kg) prolonged latency time against thermal pain generation. The latency time of AK-2a (0.5 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 4.85 ± 0.32 , 8.23 ± 0.12 , 9.45 ± 0.12 s ($P < 0.01$ vs. saline group), 10.48 ± 0.17 and 11.32 ± 0.12 s ($P < 0.001$ vs. saline group) respectively. The latency time of AK-2a (1 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 5.89 ± 0.26 , 8.55 ± 0.06 ,

Table 1 E-value (kcal/mol) and post-docking analysis of best pose of (2E,5E)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a), (1E,4E)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl) oct-1-en-3-one (AK-2a) and standard drugs with cyclooxygenase-1 (COX-1), cyclooxygenase-2 (COX-2), mu receptor, kappa receptor, delta receptor, human capsaicin receptor (HCR) and purinoceptor-3 (P2X3)

Target proteins	PDB-IDs	AK-1a			AK-2a			Standard drugs			
		E-value (kcal/mol)	No. of H-bond	Binding residues	E-value (kcal/mol)	No. of H-bond	Binding residues	Standard	E-value (kcal/mol)	No. of H-bond	Binding residues
COX 1	3N8X	-10.2	03	GLN:A:44 CYS:A:47 ARG:A:469	-9.6	02	TYR:A:130 ARG:A:469	Aspirin	-6.1	04	SER:A:126(2) GLN:A:372 GLU:B:543
COX 2	1PXX	-9.7	03	TRP:A:323 GLN:A:327 SER:B:1049	-9.6	05	GLN:C:2543 ARG:D:3044 TYR:D:3130(2) ALA-D:3156	Aspirin	-7.6	03	THR:B:1206 HIS-B:1207 TRP-B:1387
Mu receptor	5C1M	-8.7	02	ASN:A:127 HIS:A:297	-8.8	03	TYR:A:75 ASN:A:127 HIS:A:297	Morphine	-8.5	02	HIS-A:297 ASP-A:147
Kappa receptor	4DJH	-8.3	02	THR:A:63 TYR-B:313	-8.4	03	THR-B:63(2) SER-B:116	Morphine	-8.0	00	00
Delta receptor	4EJ4	-8.2	03	VAL:A:75 LYS:A:166 ASN:A:169	-8.0	02	LYS-A:108 HIS-A:278	Morphine	-7.3	00	00
Human Capsaicin receptor	3J9J	-8.8	02	ARG-C:177 GLY-C:183	-8.9	01	ARG-D:177	Capsazepine	-8.2	03	ASN-A:57 SER-A:103 TYR-A:107
P2X3	5SVL	-7.4	01	TRP-A:41	-6.9	02	LYS-A:65 PHE-A:205	Capsazepine	-5.4	02	ASP-A:266 ASN-A:279

GLN glutamine, CYS cysteine, ARG arginine, TYR tyrosine, SER serine, GLU glutamic acid, TRP tryptophan, ALA alanine, THR threonine, HIS histidine, ASN asparagine, VAL valine, LYS lysine, GLY glycine, PHE phenylalanine, ASP aspartic acid







Table 2 E-value (Kcal/mol) and post-docking analysis of best pose of (2E,5E)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a), (1E,4E)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl) oct-1-en-3-one (AK-2a) and standard drugs with cyclooxygenase-1 (COX-1), glycoprotein IIb/IIIa (GP- IIb/IIIa), glycoprotein-VI (GP-VI), purinergic receptor P₂Y₁₂ (P₂Y₁₂), prostacyclin receptor I₂ (PG-I₂) and proteinase-activated receptor 1 (PAR-1)

Target proteins	PDB-IDs	AK-1a			AK-2a			Standard drugs			
		E-value (kcal/mol)	No. of H-bond residues	Bonding residues	E-value (kcal/mol)	No. of H-bond residues	Bonding residues	Standard	E-value (kcal/mol)	No. of H-bond residues	Bonding residues
COX-1	3N8X	-10.2	03	GLN-A:44	-9.6	02	TYR-A:130	Aspirin	-6.1	04	SER-A:126(2)
				CYS-A:47			ARG-A:469				GLN-A:372
GP-IIb/IIIa	2VdM	-8.7	02	ARG-A:469	-8.4	03	ARG-A:90(2)	Tirofiban	-7.9	07	GLU-B:543
				GLN-A:18			LYS-A:124				SER-B:121
GP-VI	2G17	-7.1	02	LYS-A:124	-6.8	02	TYR-A:126,161	Hindokitol	-5.8	01	TYR-B:122
				SER-A:69(2)			00				ASP-A:159
P ₂ Y ₁₂	4PXZ	-8.5	02	LYS-A:64	-7.3	00	00	Clopidogrel	-6.0	04	PHE-A:160
				ASN-A:65			GLU-A:09				ASN-B:214
PG-I ₂	4F8K	-8.9	03	ILE-A:05	-8.4	02	GLU-A:09	Beraprost	-8.3	02	ASN-B:215(2)
				GLU-A:09			UNK-A:12				SER-A:16
PAR-1	3VW7	-10.4	02	TYR-A:187	-10.4	02	TYR-A:187,337	Vorapaxar	-12.4	06	SER-A:113(2)
				GLY-A:233			UNK-A:12				ASN-A:201(2)

GLN glutamine, CYS cysteine, ARG arginine, TYR tyrosine, SER serine, GLU glutamic acid, TRP tryptophan, ALA alanine, HIS histidine, ASN asparagine, VAL valine, LYS lysine, LEU leucine, ILE isoleucine, GLY glycine, PHE phenylalanine, ASP aspartic acid









Table 3 E-value (kcal/mol) and post-docking analysis of best pose of (2E,5E)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a), (1E,4E)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl) oct-1-en-3-one (AK-2a) and standard drugs with antithrombin-III (AT-III), factor-X (F-X), factor-II (F-II), factor-IX (F-IX) and vitamin-K epoxide reductase (VKOR)

Target proteins	PDB-IDs	AK-1a			AK-2a			Standard drugs			
		E-value (kcal/mol)	No. of H-bond	Bonding residues	E-value (kcal/mol)	No. of H-bond	Bonding residues	Standard	E-value (kcal/mol)	No. of H-bond	Bonding residues
AT-III	2B4X	-8.1	00	00	-8.2	02	ASN-I:233 ARG-I:399	Heparin SO ₄	-4.1	06	ASN-I:233 GLN-L:268(2) VAL-I:388 ARG-I:393(2)
F-II	1XMN	-8.7	01	GLY-B:223	-8.1	03	LYS-D:169 GLY-D:223 TYR-D:225	Argatroban	-8.0	08	GLU-D:39 LEU-D:40 LEU-D:41 ASN-D:143 THR-D:147B ALA-D:147C GLU-D:192
F-IX	1RFN	-9.1	02	ASN-A:48 GLY-B:114	-8.2	04	ASN-A:97 THR-A:175 SER-A:195 GLN-A:192	Pegnivacogin	-7.6	00	3D image not found
F-X	1KSN	-8.3	01	GLN-A:192	-8.2	03	TYR-A:99 SER-A:195 GLU-A:217	Apixaban	-9.2	03	TYR-A:99 GLN-A:192 SER-A:195
VKOR	3KP9	-10.3	03	ALA-A:110 MET-A:111,122	-7.6	00	00	Warfarin	-12.4	02	THR-A:34 LYS-A:41

GLN glutamine, CYS cysteine, ARG arginine, TYR tyrosine, SER serine, MET methionine, GLU glutamic acid, TRP tryptophan, ALA alanine, THR threonine, HIS histidine, ASN asparagine, VAL valine, LYS lysine, LEU leucine, ILE isoleucine, GLY glycine, PHE phenylalanine, ASP aspartic acid







10.080 \pm 0.105, 11.23 \pm 0.21 and 12.06 \pm 0.15 s (*P* < 0.001 vs. saline group) respectively. The latency time of AK-2a (3 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 6.39 \pm 0.18, 8.93 \pm 0.03 (*P* < 0.05 vs. saline group), 10.70 \pm 0.05, 11.70 \pm 0.12 and 14.49 \pm 0.25 s (*P* < 0.001 vs. saline group) respectively. The latency time of AK-2a (5 mg/kg) treated group at 0, 30, 60, 90 and 120 min were 7.96 \pm 0.15, 9.16 \pm 0.04, 11.36 \pm 0.23, 12.99 \pm 0.15 and

15.69 \pm 0.19 s (*P* < 0.001 vs. saline group) respectively as shown in Fig. 15.

Effect on AA-induced platelet aggregation inhibition

AK-1a at 1, 3, 10, 30, 100, 300 and 1000 μ M concentrations, inhibited AA-induced platelet aggregation to 2.3 \pm 0.06, 7.2 \pm 0.06, 20.4 \pm 0.06, 33.2 \pm 0.14, 55.6 \pm 0.20, 67.1 \pm 0.15 and 88.5 \pm 0.18% respectively with IC₅₀



value of 65.2 μ M. At same concentrations AK-2a inhibited AA-induced platelet aggregation to 4.3 ± 0.07 , 10.5 ± 0.09 , 28 ± 0.15 , 42.7 ± 0.22 , 62.2 ± 0.08 , 78.9 ± 0.19 and $89.8 \pm 0.13\%$ respectively with IC_{50} value of 37.7 μ M. Aspirin inhibited AA-induced platelet aggregation to 27.2 ± 0.18 , 36 ± 0.09 , 50.1 ± 0.16 , 59.7 ± 0.09 and 100% respectively with IC_{50} value of 10.01 μ M, as shown in Table 4.

Effect on ADP-induced platelet aggregation inhibition

AK-1a at 1, 3, 10, 30, 100, 300 and 1000 μ M concentrations, inhibited ADP-induced platelet aggregation to 1.81 ± 0.04 , 4.4 ± 0.04 , 13.4 ± 0.06 , 22.4 ± 0.04 , 31 ± 0.06 , 42.6 ± 0.06 and $54.1 \pm 0.06\%$ respectively with IC_{50} value of 750.4 μ M. At same concentrations AK-2a inhibited ADP-induced platelet aggregation to 4.4 ± 0.04 , 4.4 ± 0.07 , 14.2 ± 0.02 , 18.6 ± 0.06 , 30.2 ± 0.07 , 48.3 ± 0.12 and $56.8 \pm 0.06\%$ respectively with IC_{50} value of 422 μ M. Aspirin inhibited ADP-induced platelet aggregation to 3.6 ± 0.07 , 6.2 ± 0.09 , 19.1 ± 0.07 , 25 ± 0.06 , 32.8 ± 0.10 , 49.8 ± 0.12 , 49.8 ± 0.12 and $56.9 \pm 0.18\%$ respectively with IC_{50} value of 308.4 μ M, as shown in Table 4.

Effect on PRT

At 30, 100, 300 and 1000 μ M concentrations, AK-1a increased coagulation time to 137 ± 2.12 , 182.8 ± 5.59 , 224.6 ± 8.37 and 284 ± 9.46 s ($P < 0.001$ vs. saline group) respectively. AK-2a increased coagulation time to 128 ± 2.16 , 150.6 ± 2.29 , 186 ± 3.25 and 223 ± 4.47 s ($P < 0.001$ vs. saline group) respectively as shown in

Fig. 16. At 440 μ M concentration, heparin increased coagulation time to 379.40 ± 9.17 s ($P < 0.001$ vs. saline group).

Effect on BT

At 100, 300 and 1000 μ g/kg doses, AK-1a increased BT to 45.25 ± 1.75 , 59.25 ± 1.65 ($P < 0.01$ vs. saline group) and 77.75 ± 3.32 s ($P < 0.001$ vs. saline group) respectively. AK-2a increased BT to 75.25 ± 3.56 ($P < 0.01$ vs. saline group), 91.50 ± 11.11 and 120.50 ± 1.44 s ($P < 0.001$ vs. saline group) respectively as shown in Fig. 17. Heparin at 40 μ g/kg dose, increased BT to 170.75 ± 7.75 s ($P < 0.001$ vs. saline group).


Discussion

In this study, we synthesized and chemically characterized two new dibenzylidene ketone derivatives. The in-silico study carried out to get an initial information about the affinity of any compound before the start of in-vivo experiment. Docking is a preliminary tool used to check the affinity of ligands to their respective protein targets. Molecular docking has an ambient role in drug discovery and development including structure based evaluation and finding target specificity and binding affinity [25]. These interactions may exist in the form of hydrogen bonds, hydrophobic interactions and Van der Waal forces. Auto Dock Vina program was used through PyRx. It uses gradient optimization

Table 4 Inhibitory effect of (2E,5E)-2-(4-methoxybenzylidene)-5-(4-nitrobenzylidene) cyclopentanone (AK-1a) and (1E,4E)-4-(4-nitrobenzylidene)-1-(4-nitrophenyl)oct-1-en-3-one (AK-2a) and aspirin against arachidonic acid (AA) and adenosine diphosphate (ADP)-induced platelet aggregation


Treatment	Agonist	Inhibition of platelet aggregation (%) ± SEM							IC ₅₀ (μM)
		1 μM	3 μM	10 μM	30 μM	100 μM	300 μM	1000 μM	
AK-1a	AA	2.3 ± 0.06	7.2 ± 0.06	20.4 ± 0.06	33.2 ± 0.14	55.6 ± 0.20	67.1 ± 0.15	88.5 ± 0.18	65.2
	ADP	1.81 ± 0.04	4.4 ± 0.04	13.4 ± 0.06	22.4 ± 0.04	31 ± 0.06	42.6 ± 0.06	54.1 ± 0.06	750.4
AK-2a	AA	4.3 ± 0.07	10.5 ± 0.09	28 ± 0.15	42.7 ± 0.22	62.2 ± 0.08	78.9 ± 0.19	89.8 ± 0.13	37.7
	ADP	4.4 ± 0.04	4.4 ± 0.07	14.2 ± 0.02	18.6 ± 0.06	30.2 ± 0.07	48.3 ± 0.12	56.8 ± 0.06	422
Aspirin	AA	27.2 ± 0.18	36 ± 0.09	50.1 ± 0.16	59.7 ± 0.09	100 ± 0	100 ± 0	100 ± 0	10.01
	ADP	3.6 ± 0.07	6.2 ± 0.09	19.1 ± 0.07	25 ± 0.06	32.8 ± 0.10	49.8 ± 0.12	56.9 ± 0.18	308.4

Values shown as mean ± SEM, n = 4



method and it improves accuracy of binding mode predictions [26]. Hydrogen bonding is reported to be very significant in the formation of ligand protein complex [27]. Further we assessed affinity of ligands through E-value and number of hydrogen bonds against protein targets that influence analgesic, antiplatelet and anticoagulant effect. AK-1a and AK-2a showed highest binding affinity against PAR-1. AK-1a order of binding affinity against target proteins was found as: VKOR > COX-1 > COX-2 > F-IX > PG-I₂ > HCR > mu receptor > GPIIb/IIa > F-II > P₂Y₁₂ > kappa receptor > F-X > delta receptor > AT-III > P2X₃ > GP-VI. AK-2a order of binding affinity against target proteins was found as: COX-1 > COX-2 > HCR > mu receptor > kappa receptor > GPIIb/IIIa > PG-I₂ > AT-III > F-IX > F-X > F-II > delta

receptor > VKOR > P₂Y₁₂ > P₂X₃ > GPVI. We can infer that our compounds have analgesic, antiplatelet and anticoagulant actions. The analgesic activity was studied using two standard protocols i.e. acetic acid induced writhing method and hot plate assay to evaluate the peripheral and central effects of analgesia [28]. Basically writhing is an abdominal constriction caused by the release of different types of mediators after the i.p injection of acetic acid. This noxious response can be prevented by drugs which have the ability to stop the synthesis of these chemicals. The reduction in the number of writhes in treated group explains the same phenomenon of blocking the production of mediators by inhibiting COX-2 by the test compounds. Analgesic actions of AK-1a and AK-2a are proposed as inhibition of prostanoid release from cyclooxygenase involved in visceral nociception induced by acetic acid [29]. The central nociceptive effects were validated through hotplate assay [30]. AK-1a and AK-2a showed dose-dependent analgesic response, while AK-2a is found to be potent, as dose ≥ 10 mg/kg cannot be used for the analgesic activity. Significant response against acetic acid-induced writhing and hotplate assay by AK-1a and AK-2a explains central as well as peripheral activity of dibenzylidene ketone derivatives [31]. In acetic acid-induced writhing at higher dose AK-2a showed significant response, it can be further checked for anti-inflammatory response. The nociceptive behavior in the acetic acid-induced writhing test occurs due to synthesis of pain mediators including prostaglandins due to induction of COX-2 that results increased in pain sensitivity after acetic acid injection [32, 33]. Acetic acid produces nociception by releasing chemical mediators such as serotonin, histamine, prostaglandins, bradykinins and substance P due to induction of COX-2 that results in increased pain sensitivity after acetic acid injection. The acetic-induced writhing test is also



sensitive to adrenoceptor agonists and opioid agonists which through appropriate receptor stimulation in the peritoneal cavity cause reduction in pain perception [34, 35]. This test involves both central and peripheral mechanisms in the early phase of the test [36]. However, hot plate test is regarded as a suitable model for the involvement of central mechanisms [37, 38]. PAR-1 activation leads to stimulation of arachidonic acid release and thrombin signaling. Arachidonic acid enhances the activation of platelet aggregation cascade [39, 40]. This can be a proposed mechanism of action for AK-1a and AK-2a as antiplatelet and anticoagulant agents. As per computational study results, both can be a potential antagonist of PAR-1 which was further validated. Curcumin analogues inhibit platelet aggregation and repress thrombosis. Dibenzylidene ketone derivatives used in this study, having ketone moiety showed significant antiplatelet and anticoagulant response [41] presence of methoxy group in AK-1a enhanced its biological activity [42]. Previous studies revealed role of curcumin derivatives as a vitamin K antagonist [43], so as these dibenzylidene ketone derivatives. Anticoagulant actions of AK-1a and AK-2a were also validated by the presence of hydrophobic groups [44].

Conclusions

The present study reports the newly synthesized dibenzylidene ketone derivatives AK-1a and AK-2a showed high binding affinities against different protein targets involved in mediation of pain, platelet aggregation and blood coagulation process. The pharmacological investigations based on in-silico, in-vitro and in-vivo studies revealed their analgesic, antiplatelet and anticoagulant

actions. These are promising findings, since the production of dibenzylidene compounds is a simple, cheap and feasible process.

Authors' contributions

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication. TA carried out the computational studies, experimental work, analyzed the data and documentation. AK and FA supervised the research project. EF and AK synthesized dibenzylidene ketone derivatives. MA and ZD revised the final manuscript. All authors read and approved the final manuscript.

Author details

¹ Riphah Institute of Pharmaceutical Sciences, Riphah International University, Islamabad, Pakistan. ² LaBioMMi, Department of Chemistry, Federal University of São Carlos, CP 676, São Carlos, SP 13565-905, Brazil. ³ Department of Chemistry, Woman University Swabi, GulooDehri, Topi Road, Swabi, KP 23340, Pakistan. ⁴ Basic Sciences Department, College of Science and Health Professions-(COSHP-J), King Saud bin Abdulaziz University for Health Sciences, Jeddah, Saudi Arabia. ⁵ Present Address: Department of Pharmacy, Capital University of Science and Technology, Islamabad, Pakistan.

Acknowledgements

The authors are thankful to Riphah Academy of Research and Education, Riphah International University, for partial financial support of the study.

Competing interests

The authors declare that they have no competing interests.

Funding

There are no specific funding for the study.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 9 March 2018 Accepted: 29 November 2018

Published online: 06 December 2018

References

- Loeser JD, Melzack R (1999) Pain: an overview. *Lancet* 353(9164):1607–1609. [https://doi.org/10.1016/S0140-6736\(99\)01311-2](https://doi.org/10.1016/S0140-6736(99)01311-2)
- Zulfiker A, Rahman MM, Hossain MK, Hamid K, Mazumder M, Rana MS (2010) In vivo analgesic activity of ethanolic extracts of two medicinal plants-*Scoparia dulcis* L. and *Ficus racemosa* Linn. *Biol Med* 2:42–48
- O'Brien J (1995) Platelet activation at high and low shear is followed by inactivation: the clinical relevance. *Platelets* 6:242–243. <https://doi.org/10.3109/09537109509078461>
- Warfarin Antiplatelet Vascular Evaluation Trial Investigators (2007) Oral anticoagulant and antiplatelet therapy and peripheral arterial disease. *N Engl J Med* 2007:217–227. <https://doi.org/10.1056/NEJMoa065959>
- Mackman N (2008) Triggers, targets and treatments for thrombosis. *Nature* 451:914. <https://doi.org/10.1038/nature06797>
- Dahlbäck B (2000) Blood coagulation. *Lancet* 355:1627–1632. [https://doi.org/10.1016/S0140-6736\(00\)02225-X](https://doi.org/10.1016/S0140-6736(00)02225-X)
- Maynard J, Heckman C, Pitlick F, Nemerson Y (1975) Association of tissue factor activity with the surface of cultured cells. *J Clin Invest* 55:814. <https://doi.org/10.1172/JCI107992>
- Davie EW, Fujikawa K, Kisiel W (1991) The coagulation cascade: initiation, maintenance, and regulation. *Biochem* 30:10363–10370. <https://doi.org/10.1021/bi00107a001>
- Schlitt A, Rubboli A, Airaksinen KEJ, Lip GY (2013) Antiplatelet therapy and anticoagulants. *Lancet* 382(9886):24–25
- Shen L, Ji H-F (2007) Theoretical study on physicochemical properties of curcumin. *Spectrochim Acta Part A Mol Biomol Spectrosc* 67:619–623. <https://doi.org/10.1016/j.saa.2006.08.018>

- Hatcher H, Planalp R, Cho J, Torti F, Torti S (2008) Curcumin: from ancient medicine to current clinical trials. *Cell Mol Life Sci* 65:1631–1652. <https://doi.org/10.1007/s00018-008-7452-4>
- Srivastava R, Dikshit M, Srimal R, Dhawan B (1985) Anti-thrombotic effect of curcumin. *Thromb Res* 40:413–417
- Srivastava K (1989) Extracts from two frequently consumed spices—cumin (*Cuminum cyminum*) and turmeric (*Curcuma longa*)—inhibit platelet aggregation and alter eicosanoid biosynthesis in human blood platelets. *Prostaglandins Leukot Essent Fatty Acids* 37:57–64. [https://doi.org/10.1016/0952-3278\(89\)90187-7](https://doi.org/10.1016/0952-3278(89)90187-7)
- Dinkova-Kostova AT, Massiah MA, Bozak RE, Hicks RJ, Talalay P (2001) Potency of Michael reaction acceptors as inducers of enzymes that protect against carcinogenesis depends on their reactivity with sulfhydryl groups. *Proc Natl Acad Sci* 98(6):3404–3409. <https://doi.org/10.1073/pnas.051632198>
- Sahu PK, Sahu PK, Gupta S, Thavaselvam D, Agarwal D (2012) Synthesis and evaluation of antimicrobial activity of 4H-pyrimido[2,1-b]benzothiazole, pyrazole and benzylidene derivatives of curcumin. *Eur J Med Chem* 54:366–378. <https://doi.org/10.1016/j.ejmech.2012.05.020>
- Din ZU, Lazarin-Bidóia D, Kaplum V, Garcia FP, Nakamura CV, Rodrigues-Filho E (2016) The structure design of biotransformed unsymmetrical nitro-contained 1,5-diaryl-3-oxo-1,4-pentadienyls for the anti-parasitic activities. *J Chem, Arab*. <https://doi.org/10.1016/j.arabjc.2016.03.005>
- Trott O, Olson AJ (2010) AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* 31:455–461. <https://doi.org/10.1002/jcc.21334>
- Dallakyan S, Olson AJ (2015) Small-molecule library screening by docking with pyrx. *Chemical Biology: Methods Protocols*. Humana Press, New York, pp 243–250. <https://doi.org/10.1007/978-1-4939-2269-7>
- Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK, Olson AJ (1998) Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function. *J Comput Chem* 19:1639–1662
- Major C, Pleuvry BJ (1971) Effects of α -methyl-*p*-tyrosine, *p*-chlorophenylalanine, l - β -(3,4-dihydroxyphenyl) alanine, 5-hydroxytryptophan and diethylthiocarbamate on the analgesic activity of morphine and methylamphetamine in the mouse. *Br J Pharmacol* 42:512–521. <https://doi.org/10.1111/j.1476-5381.1971.tb07137.x>
- Adzu B, Amos S, Kapu S, Gamanil K (2003) Anti-inflammatory and anti-nociceptive effects of *Sphaeranthus senegalensis*. *J Ethnopharmacol* 84(2):169–173. [https://doi.org/10.1016/S0378-8741\(02\)00295-7](https://doi.org/10.1016/S0378-8741(02)00295-7)
- Bauriedel G, Skowasch D, Schneider M, Andrié R, Jabs A, Lüderitz B (2003) Antiplatelet effects of angiotensin-converting enzyme inhibitors compared with aspirin and clopidogrel: a pilot study with whole-blood aggregometry. *Am Heart J* 145:343–348. <https://doi.org/10.1067/mhj.2003.22>
- Dey P, Bhakta T (2012) Evaluation of in vitro anticoagulant activity of *Molineria recurvata* leaf extract. *J Nat Prod Plant Resour* 2:685–688
- Gangaraju S, Manjappa B, Subbaiah GK, Kempaiah K, Shinde M, Sananigaiah D (2015) Anticoagulant and antiplatelet activities of jackfruit (*Artocarpus heterophyllus*) seed extract. *Int J Pharm Pharm Sci* 7:187–191. <https://doi.org/10.5530/pj.2015.3.5>
- Morris GM, Lim-Wilby M (2008) Molecular docking. *Molecular modeling of proteins*. Humana Press, New York, pp 365–382
- Ayyapa B, Kanchi S, Singh P, Sabela MI, Dovey M, Bisetty K (2015) Analytical evaluation of steviol glycosides by capillary electrophoresis supported with molecular docking studies. *JICS* 12(1):127–136. <https://doi.org/10.1007/s13738-014-0465-z>
- Wang R, Lai L, Wang S (2002) Further development and validation of empirical scoring functions for structure-based binding affinity prediction. *J Comput Aided Mol Des* 16:11–26
- Gené RM, Segura L, Adzet T, Marin E, Iglesias J (1998) Heterothecainuloides: anti-inflammatory and analgesic effect. *J Ethnopharmacol* 60(2):157–162
- Ulugöl A, Özyigit F, Yesilyurt Ö, Dogrul A (2006) The additive antinociceptive interaction between win 55,212-2, a cannabinoid agonist, and ketorolac. *Anesth Analg* 102:443–447. <https://doi.org/10.1213/01.ane.0000194587.94260.1d>
- Arslan R, Bektas N, Ozturk Y (2010) Antinociceptive activity of methanol extract of fruits of *Capparis ovata* in mice. *J Ethnopharmacol* 131:28–32. <https://doi.org/10.1016/j.jep.2010.05.060>
- Chapman CR, Casey K, Dubner R, Foley K, Gracely R, Reading A (1985) Pain measurement: an overview. *Pain* 22:1–31. [https://doi.org/10.1016/0304-3959\(85\)90145-9](https://doi.org/10.1016/0304-3959(85)90145-9)
- Matsumoto H, Naraba H, Ueno A et al (1998) Induction of cyclooxygenase-2 causes an enhancement of writhing response in mice. *Eur J Pharmacol* 352:47–52. [https://doi.org/10.1016/S0014-2999\(98\)00340-9](https://doi.org/10.1016/S0014-2999(98)00340-9)
- Ballou LR, Botting RM, Goorha S, Zhang J, Vane JR (2000) Nociception in cyclooxygenase isozyme-deficient mice. *Proc Natl Acad Sci* 97:10272–10276. <https://doi.org/10.1073/pnas.180319297>
- Bentley GA, Newton SH, Starr J (1981) Evidence for an action of morphine and enkephalins on sensory nerves endings in the mouse peritoneum. *Br J Pharmacol* 73:325–332. <https://doi.org/10.1111/j.1476-5381.1981.tb10425.x>
- Bentley GA, Newton SH, Starr J (1983) Studies on the antinociceptive action of α -agonist drugs and their interaction with opioid mechanisms. *Br J Pharmacol* 73:125–134. <https://doi.org/10.1111/j.1476-5381.1983.tb10504.x>
- Chen TF, Tsai HY, Tian-Shang W (1995) Anti-inflammatory and analgesic activities from the roots of *Angelica pubescens*. *Planta Med* 61:2–8. <https://doi.org/10.1055/s-2006-957987>
- Pini LA, Vitale G, Ottani A, Sandrini M (1997) Naloxone-reversible antinociception by paracetamol in the rat. *J Pharmacol Exp Ther* 280:934–940
- Hosseinzadeh H, Younesi HM (2002) Antinociceptive and anti-inflammatory effects of *Crocus sativus* stigma and petal extracts in mice. *BMC Pharmacol* 2:7. <https://doi.org/10.1186/1471-2210-2-7>
- Coughlin SR (2000) Thrombin signalling and protease-activated receptors. *Nature* 407:258–264. <https://doi.org/10.1038/35025229>
- Howat JM, Creer MH, Rickard A (2001) Stimulation of protease activated receptors on RT4 cells mediates arachidonic acid release via Ca^{2+} independent phospholipase A_2 . *J Urol* 165:2063–2067. [https://doi.org/10.1016/S0022-5347\(05\)66295-7](https://doi.org/10.1016/S0022-5347(05)66295-7)
- Bukhari SNA, Jantan IB, Jasamai M, Ahmad W, Amjad MWB (2013) Synthesis and biological evaluation of curcumin analogues. *J Med Sci* 13:501–513. <https://doi.org/10.3923/jms.2013.501.513>
- Cao Y-K, Li H-J, Song Z-F, Li Y, Huai Q-Y (2014) Synthesis and biological evaluation of novel curcuminoid derivatives. *Molecules* 19:16349–16372. <https://doi.org/10.3390/molecules191016349>
- Hoult J, Paya M (1996) Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Vasc Pharmacol* 27:713–722. [https://doi.org/10.1016/0306-3623\(95\)02112-4](https://doi.org/10.1016/0306-3623(95)02112-4)
- Manikandan P, Sumitra M, Aishwarya S, Manohar BM, Lokanadam B, Puvanakrishnan R (2004) Curcumin modulates free radical quenching in myocardial ischaemia in rats. *Int J Biochem Cell Biol* 36:1967–1980. <https://doi.org/10.1016/j.biocel.2004.01.030>