

SOME RESULTS ON DEGENERATE FUBINI AND DEGENERATE BELL POLYNOMIALS

Taekyun Kim and Dae San Kim*

The aim of this paper is to further study some properties and identities on the degenerate Fubini and the degenerate Bell polynomials which are degenerate versions of the Fubini and the Bell polynomials, respectively. Especially, we find several expressions for the generating function of the sum of the values of the generalized falling factorials at positive consecutive integers.

1. INTRODUCTION

Carlitz [3] initiated a study of degenerate versions of Bernoulli and Euler polynomials, namely the degenerate Bernoulli and the degenerate Euler polynomials. In recent years, some mathematicians have explored degenerate versions of many special polynomials and numbers, which include the degenerate Bernoulli numbers of the second kind, the degenerate Stirling numbers of the first and second kinds, the degenerate Bell numbers and polynomials, and so on (see [6,8,10-12,14-17,19] and the references therein). Many interesting results on such degenerate versions together with their applications were found by employing several different tools such as combinatorial methods, generating functions, p -adic analysis, umbral calculus techniques, differential equations, probability theory, operator theory, special functions and analytic number theory. It is remarkable that the exploration for degenerate versions is not only limited to polynomials and numbers but also extended to transcendental functions like gamma functions (see [13]).

Recently, the degenerate Fubini polynomials and the degenerate Bell polynomials are introduced (see [12,16]). The aim of this paper is to further study some properties and identities on the degenerate Fubini and the degenerate Bell

*Corresponding author. Taekyun Kim

2020 Mathematics Subject Classification. 11B73, 11B83

Keywords and Phrases. Degenerate Fubini polynomials, Degenerate Bell polynomials

polynomials which are degenerate versions of the Fubini and the Bell polynomials, respectively. Especially, we find several expressions for the generating functions of the sum $S_{n,\lambda}(p) = (1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda}$ of the values of the generalized falling factorials at positive consecutive integers. The outline of this paper is as follows. In Section 1, we recall degenerate exponential functions, degenerate logarithms, degenerate Stirling numbers of the first kind and degenerate Stirling numbers of the second kind which are respectively degenerate versions of exponentials, logarithms, Stirling numbers of the first kind and Stirling numbers of the second kind. Also, we recall the degenerate Bell polynomials, the degenerate polylogarithm and the degenerate Fubini polynomials. In Section 2, we derive recurrence relations for the degenerate Bell polynomials in Theorems 1 and 2. In Theorem 4 and 7, the exponential generating function of the sum $S_{n,\lambda}(p)$ is determined in terms of the degenerate Bell polynomials. Another exponential generating function for the sum $S_{n,\lambda}(p)$ is obtained in Theorem 6 in terms of the degenerate Stirling numbers of the second kind. In Theorem 8, the generating function for $S_{n,\lambda}(p)$ is determined in terms of the degenerate Fubini polynomial. In Theorem 12, obtained is a differential equation satisfied by the degenerate Fubini polynomials. Finally, we conclude our paper in Section 3.

The generalized falling factorial sequence is defined by

$$(1) \quad (x)_{0,\lambda} = 1, \quad (x)_{n,\lambda} = x(x-\lambda)(x-2\lambda) \cdots (x-(n-1)\lambda), \quad (n \geq 1),$$

where λ is a real number (see [8,12]).

For any $\lambda \in \mathbb{R}$, the degenerate exponentials are defined by

$$(2) \quad e_{\lambda}^x(t) = \sum_{n=0}^{\infty} (x)_{n,\lambda} \frac{t^n}{n!}, \quad (\text{see [11, 14]}).$$

In particular, for $x = 1$, $e_{\lambda}(t) = e_{\lambda}^1(t) = \sum_{n=0}^{\infty} (1)_{n,\lambda} \frac{t^n}{n!}$. Note that $\lim_{\lambda \rightarrow 0} e_{\lambda}^x(t) = e^{xt}$.

The degenerate Bell polynomials are considered by Kim-Kim and given by

$$(3) \quad e^{x(e_{\lambda}(t)-1)} = \sum_{n=0}^{\infty} \phi_{n,\lambda}(x) \frac{t^n}{n!}, \quad (\text{see [14, 15]}).$$

From (1), (2) and (3) we note that

$$\lim_{n \rightarrow \infty} \phi_{n,\lambda}(x) = \phi_n(x), \quad (n \geq 0),$$

where $\phi_n(x)$ are the ordinary Bell polynomials given by

$$(4) \quad e^{x(e^t-1)} = \sum_{n=0}^{\infty} \phi_n(x) \frac{t^n}{n!}, \quad (\text{see [6-12, 14-20]}).$$

In [8], the degenerate Stirling numbers of the first kind are defined by

$$(5) \quad (x)_n = \sum_{k=0}^n S_{1,\lambda}(n, k)(x)_{k,\lambda}, \quad (n \geq 0),$$

where $(x)_0 = 1$, $(x)_n = x(x-1) \cdots (x-n+1)$, $(n \geq 1)$.

Note that $\lim_{\lambda \rightarrow 0} S_{1,\lambda}(n, k) = S_1(n, k)$ are the ordinary Stirling number of the first kind given by

$$(x)_n = \sum_{k=0}^n S_1(n, k) x^k, \quad (n \geq 0), \quad (\text{see [1-5, 7]}).$$

As the inversion formula of (5), the degenerate Stirling numbers of the second kind are introduced by Kim-Kim and given by

$$(6) \quad (x)_{n,\lambda} = \sum_{k=0}^n S_{2,\lambda}(n, k) (x)_k, \quad (n \geq 0), \quad (\text{see [8]}).$$

Let $\log_\lambda t$ be the compositional inverse of $e_\lambda(t)$. Then we have

$$(7) \quad \log_\lambda(1+t) = \sum_{n=1}^{\infty} \frac{\lambda^{n-1} (1)_{n,1/\lambda}}{n!} t^n, \quad (\text{see [8]}).$$

Note that $\lim_{\lambda \rightarrow 0} \log_\lambda(1+t) = \log(1+t)$.

For $k \in \mathbb{Z}$, the degenerate polylogarithm function is defined by Kim-Kim and given by

$$(8) \quad \text{Li}_{k,\lambda}(t) = \sum_{n=1}^{\infty} \frac{(-\lambda)^{n-1} (1)_{n,1/\lambda}}{(n-1)! n^k} t^n, \quad (\text{see [8]}).$$

Note that $\lim_{\lambda \rightarrow 0} \text{Li}_{k,\lambda}(t) = \text{Li}_k(t) = \sum_{n=1}^{\infty} \frac{x^n}{n^k}$ is the polylogarithm function and $\text{Li}_{1,\lambda}(t) = -\log_\lambda(1-t)$.

From (5) and (10), we can derive the generating functions of degenerate Stirling numbers which are given by

$$(9) \quad \frac{1}{k!} (\log_\lambda(1+t))^k = \sum_{n=k}^{\infty} S_{1,\lambda}(n, k) \frac{t^n}{n!},$$

and

$$(10) \quad \frac{1}{k!} (e_\lambda(t) - 1)^k = \sum_{n=k}^{\infty} S_{2,\lambda}(n, k) \frac{t^n}{n!}, \quad (k \geq 0), \quad (\text{see [8]}).$$

In [15], the degenerate Fubini polynomials are defined by

$$(11) \quad \frac{1}{1 - x(e_\lambda(t) - 1)} = \sum_{n=0}^{\infty} F_{n,\lambda}(x) \frac{t^n}{n!}.$$

Thus, by (11), we get

$$(12) \quad F_{n,\lambda}(x) = \sum_{k=0}^n k! S_{2,\lambda}(n, k) x^k, \quad (n \geq 0), \quad (\text{see [15]}).$$

By (3) and (6) we easily get

$$(13) \quad \begin{aligned} \phi_{n,\lambda}(x) &= \sum_{k=0}^n S_{2,\lambda}(n, k) x^k, \quad (n \geq 0), \\ &= e^{-x} \sum_{k=0}^{\infty} \frac{(k)_{n,\lambda}}{k!} x^k, \quad (\text{see [9, 10, 14, 15]}). \end{aligned}$$

2. SOME NEW FORMULAE ON DEGENERATE FUBINI AND BELL POLYNOMIALS

First, by (3), (4), (5) and (6) we get

$$(14) \quad \begin{aligned} \frac{d}{dx} e^{x(e_\lambda(t)-1)} &= (e_\lambda(t) - 1) e^{x(e_\lambda(t)-1)} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} (1)_{n-k,\lambda} \phi_{k,\lambda}(x) - \phi_{n,\lambda}(x) \right) \frac{t^n}{n!}. \end{aligned}$$

Thus, by (3) and (14), we get

$$(15) \quad \begin{aligned} \phi'_{n,\lambda}(x) &= \frac{d}{dx} \phi_{n,\lambda}(x) = \sum_{k=0}^n \binom{n}{k} (1)_{n-k,\lambda} \phi_{k,\lambda}(x) - \phi_{n,\lambda}(x) \\ &= \sum_{k=0}^{n-1} \binom{n}{k} (1)_{n-k,\lambda} \phi_{k,\lambda}(x), \quad (n \geq 1). \end{aligned}$$

From (3), we note that

$$(16) \quad \begin{aligned} \sum_{n=0}^{\infty} \phi_{n+1,\lambda}(x) \frac{t^n}{n!} &= \frac{d}{dt} e^{x(e_\lambda(t)-1)} = x e_\lambda^{1-\lambda}(t) e^{x(e_\lambda(t)-1)} \\ &= x \sum_{l=0}^{\infty} (1-\lambda)_{l,\lambda} \frac{t^l}{l!} \sum_{k=0}^{\infty} \phi_{k,\lambda}(x) \frac{t^k}{k!} \\ &= \sum_{n=0}^{\infty} \left(x \sum_{k=0}^n \binom{n}{k} \phi_{k,\lambda}(x) (1-\lambda)_{n-k,\lambda} \right) \frac{t^n}{n!}. \end{aligned}$$

By comparing the coefficients on both sides of (15), we have

$$(17) \quad \phi_{n+1,\lambda}(x) = x \sum_{k=0}^n \binom{n}{k} \phi_{k,\lambda}(x) (1-\lambda)_{n-k,\lambda}, \quad (n \geq 0).$$

From the definition in (1), we easily get

$$(18) \quad (1 - \lambda)_{n-k, \lambda} = (1)_{n-k, \lambda} (1 - (n - k)\lambda).$$

From (17) and (18), we have

$$(19) \quad \begin{aligned} \phi_{n+1, \lambda}(x) &= x \sum_{k=0}^n \binom{n}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda} (1 - (n - k)\lambda) \\ &= x \sum_{k=0}^n \binom{n}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda} - x\lambda \sum_{k=0}^n \binom{n}{k} (n - k) (1)_{n-k, \lambda} \phi_{k, \lambda}(x) \\ &= x \sum_{k=0}^n \binom{n}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda} - nx\lambda \sum_{k=0}^n \binom{n-1}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda}. \end{aligned}$$

Therefore, by (19), we obtain the following theorem.

Theorem 1. For $n \in \mathbb{Z}$ with $n \geq 0$, we have

$$\sum_{k=0}^n \binom{n}{k} \phi_{n, k}(x) (1)_{n-k, \lambda} = \frac{1}{x} \phi_{n+1, \lambda}(x) + n\lambda \sum_{k=0}^{n-1} \binom{n-1}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda}.$$

By (15) and Theorem 1, we obtain the following theorem.

Theorem 2. For $n \in \mathbb{Z}$ with $n \geq 0$, we have

$$\frac{1}{x} \phi_{n+1, \lambda}(x) = \phi'_{n, \lambda}(x) + \phi_{n, \lambda}(x) - n\lambda \sum_{k=0}^{n-1} \binom{n-1}{k} \phi_{k, \lambda}(x) (1)_{n-k, \lambda}.$$

Let

$$(20) \quad \begin{aligned} Y_{\lambda}(x, p) &= \sum_{n=1}^{\infty} \sum_{k=1}^n (k)_{p, \lambda} \frac{x^n}{n!} \\ &= \sum_{n=1}^{\infty} \left((1)_{p, \lambda} + (2)_{p, \lambda} + \cdots + (n)_{p, \lambda} \right) \frac{x^n}{n!} \end{aligned}$$

where p is a positive integer.

From (20), we note that

$$(21) \quad Y_{\lambda}(x, p) = \sum_{k=1}^{\infty} (k)_{p, \lambda} \sum_{n=k}^{\infty} \frac{x^n}{n!} = \sum_{k=1}^{\infty} (k)_{p, \lambda} \left(e^x - \sum_{l=0}^{k-1} \frac{x^l}{l!} \right).$$

In (13), we note that

$$(22) \quad e^x \phi_{p, \lambda}(x) = \sum_{n=1}^{\infty} \frac{(n)_{p, \lambda}}{n!} x^n, \quad (p \in \mathbb{Z}).$$

Let

$$(23) \quad \begin{aligned} y_{p,\lambda}(x) &= Y_\lambda(x, p) - e^x \phi_{p,\lambda}(x) \\ &= \sum_{n=2}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n-1)_{p,\lambda} \right) \frac{x^n}{n!}, \quad (p \in \mathbb{N}). \end{aligned}$$

By (21), (22) and (23), we get

$$(24) \quad \begin{aligned} y'_{p,\lambda}(x) &= \frac{d}{dx} y_{p,\lambda}(x) = \sum_{n=2}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n-1)_{p,\lambda} \right) \frac{x^{n-1}}{(n-1)!} \\ &= \sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) \frac{x^n}{n!} \\ &= \sum_{n=1}^{\infty} \left((1)_{p,\lambda} + \cdots + (n-1)_{p,\lambda} \right) \frac{x^n}{n!} + \sum_{n=1}^{\infty} (n)_{p,\lambda} \frac{x^n}{n!} \\ &= y_{p,\lambda}(x) + \sum_{n=1}^{\infty} (n)_{p,\lambda} \frac{x^n}{n!} = y_{p,\lambda}(x) + e^x \phi_{p,\lambda}(x). \end{aligned}$$

From (24), we have the following differential equation.

Proposition 3. *For $p \in \mathbb{N}$, we have*

$$y'_{p,\lambda}(x) - y_{p,\lambda}(x) = e^x \phi_{p,\lambda}(x).$$

By Proposition 3, we easily get

$$(25) \quad \frac{d}{dx} \left(e^{-x} y_{p,\lambda}(x) \right) = \phi_{p,\lambda}(x).$$

Thus, by (25), we get

$$(26) \quad y_{p,\lambda}(x) = e^x \int_0^x \phi_{p,\lambda}(t) dt.$$

Therefore, by (20), (23) and (26), we obtain the following theorem.

Theorem 4. *For $p \in \mathbb{N}$, we have*

$$\sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) \frac{t^n}{n!} = e^x \phi_{p,\lambda}(x) + e^x \int_0^x \phi_{p,\lambda}(t) dt.$$

In [10], considered are the degenerate differential operators which are given by

$$(27) \quad \left(x \frac{d}{dx} \right)_{n,\lambda} = \left(x \frac{d}{dx} \right) \left(x \frac{d}{dx} - \lambda \right) \cdots \left(x \frac{d}{dx} - (n-1)\lambda \right), \quad (n \geq 1).$$

For $p \in \mathbb{N}$, by (27), we get

$$(28) \quad \left(x \frac{d}{dx}\right)_{p,\lambda} e^x = \sum_{n=1}^{\infty} \frac{(n)_{p,\lambda}}{n!} x^n = e^x \phi_{p,\lambda}(x).$$

Therefore, by (28), we obtain the following theorem.

Theorem 5. *For $p \in \mathbb{N}$, we have*

$$\left(x \frac{d}{dx}\right)_{p,\lambda} e^x = \sum_{n=1}^{\infty} \frac{(n)_{p,\lambda}}{n!} x^n = e^x \phi_{p,\lambda}(x).$$

From (23) and (26), we note that

$$(29) \quad \begin{aligned} Y_\lambda(x, p) &= e^x \phi_{p,\lambda}(x) + e^x \int_0^x \phi_{p,\lambda}(t) dt \\ &= e^x \phi_{p,\lambda}(x) + e^x \sum_{k=0}^p S_{2,\lambda}(p, k) \int_0^x t^k dt \\ &= e^x \phi_{p,\lambda}(x) + e^x \sum_{k=0}^p S_{2,\lambda}(p, k) \frac{x^{k+1}}{k+1} \\ &= e^x \sum_{k=0}^p \left(x^k + \frac{x^{k+1}}{k+1}\right) S_{2,\lambda}(p, k). \end{aligned}$$

Therefore, by (20) and (29), we obtain the following theorem.

Theorem 6. *For $p \in \mathbb{N}$, we have*

$$\sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda}\right) \frac{x^n}{n!} = e^x \sum_{k=0}^p \left(x^k + \frac{x^{k+1}}{k+1}\right) S_{2,\lambda}(p, k).$$

By (6), we easily get

$$(30) \quad S_{2,\lambda}(n+1, k) = S_{2,\lambda}(n, k-1) + (k - n\lambda) S_{2,\lambda}(n, k),$$

where $n, k \geq 0$ with $n \geq k$.

From (27), (28), (30) and noting $S_{2,\lambda}(p, k) = 0$ for $p \geq 1$, we note that

(31)

$$\begin{aligned}
Y_\lambda(x, p) &= e^x \sum_{k=0}^p \left(x^k + \frac{x^{k+1}}{k+1} \right) S_{2,\lambda}(p, k) \\
&= e^x \sum_{k=1}^{p+1} x^k S_{2,\lambda}(p, k) + e^x \sum_{k=0}^p \frac{x^{k+1}}{k+1} S_{2,\lambda}(p, k) \\
&= e^x \sum_{k=1}^{p+1} \left(S_{2,\lambda}(p, k)(k - p\lambda) + S_{2,\lambda}(p, k-1) \right) \frac{x^k}{k} + p\lambda e^x \sum_{k=1}^{p+1} \frac{S_{2,\lambda}(p, k)}{k} x^k \\
&= e^x \sum_{k=1}^{p+1} \frac{1}{k} S_{2,\lambda}(p+1, k) x^k + p\lambda e^x \sum_{k=1}^{p+1} \frac{1}{k} S_{2,\lambda}(p, k) x^k \\
&= e^x \sum_{k=1}^{p+1} \left(\frac{S_{2,\lambda}(p+1, k)}{k} + \frac{p\lambda}{k} S_{2,\lambda}(p, k) \right) x^k \\
&= e^x \int_0^x \left(\phi_{p+1,\lambda}(t) + p\lambda \phi_{p,\lambda}(t) \right) \frac{dt}{t},
\end{aligned}$$

where $p \in \mathbb{N}$ and $\frac{1}{t} \phi_{p+1,\lambda}(t) = \sum_{k=1}^{p+1} S_{2,\lambda}(p+1, k) t^{k-1}$.

Therefore, by (31), we obtain the following theorem.

Theorem 7. *For $p \in \mathbb{Z}$, we have*

$$\sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) \frac{x^n}{n!} = e^x \int_0^x \left(\phi_{p+1,\lambda}(t) + p\lambda \phi_{p,\lambda}(t) \right) \frac{dt}{t}.$$

From (12), we note that

(32)

$$\begin{aligned}
\sum_{l=0}^{\infty} (l)_{p,\lambda} x^l &= \sum_{l=0}^{\infty} x^l \sum_{k=0}^p S_{2,\lambda}(p, k) (l)_k = \sum_{l=0}^{\infty} x^l \sum_{k=0}^p S_{2,\lambda}(p, k) \binom{l}{k} k! \\
&= \sum_{l=k}^{\infty} x^l \sum_{k=0}^p S_{2,\lambda}(p, k) \binom{l}{k} k! = \sum_{k=0}^p S_{2,\lambda}(p, k) k! \sum_{l=0}^{\infty} \binom{l+k}{k} x^{l+k} \\
&= \sum_{k=0}^p S_{2,\lambda}(p, k) k! x^k \sum_{l=0}^{\infty} \binom{l+k}{k} x^l = \sum_{k=0}^p S_{2,\lambda}(p, k) k! x^k \left(\frac{1}{1-x} \right)^{k+1} \\
&= \frac{1}{1-x} \sum_{k=0}^p S_{2,\lambda}(p, k) k! \left(\frac{x}{1-x} \right)^k = \frac{1}{1-x} F_{p,\lambda} \left(\frac{x}{1-x} \right).
\end{aligned}$$

For $p \in \mathbb{Z}$ with $p \geq 0$, and by using (32), we have

$$(33) \quad \left(x \frac{d}{dx}\right)_{p,\lambda} \frac{1}{1-x} = \sum_{n=0}^{\infty} \left(x \frac{d}{dx}\right)_{p,\lambda} x^n = \sum_{n=0}^{\infty} (n)_{p,\lambda} x^n = \frac{1}{1-x} F_{p,\lambda} \left(\frac{x}{1-x}\right).$$

Thus, for $p \in \mathbb{N}$, we have

$$(34) \quad \begin{aligned} \sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) x^n &= \sum_{n=1}^{\infty} x^n \sum_{k=1}^n (k)_{p,\lambda} \\ &= \sum_{k=1}^{\infty} (k)_{p,\lambda} \sum_{n=k}^{\infty} x^n = \frac{1}{1-x} \sum_{k=1}^{\infty} (k)_{p,\lambda} x^k = \frac{1}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right). \end{aligned}$$

Therefore, by (33) and (34), we obtain the following theorem.

Theorem 8. *For $p \geq 0$, we have*

$$\left(x \frac{d}{dx}\right)_{p,\lambda} \frac{1}{1-x} = \sum_{n=0}^{\infty} (n)_{p,\lambda} x^n = \frac{1}{1-x} F_{p,\lambda} \left(\frac{x}{1-x}\right).$$

In addition, for $p \in \mathbb{N}$, we have

$$\sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) x^n = \frac{1}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right).$$

Now, we observe that, for $p \in \mathbb{N}$,

$$(35) \quad \begin{aligned} \sum_{n=1}^{\infty} (n)_{p,\lambda} \left(\frac{1}{1-x} - \sum_{l=0}^n x^l \right) &= \sum_{n=1}^{\infty} (n)_{p,\lambda} \frac{x^{n+1}}{1-x} \\ &= \frac{x}{1-x} \sum_{n=1}^{\infty} (n)_{p,\lambda} x^n = \frac{x}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right). \end{aligned}$$

Therefore, by (35), we obtain the following corollary.

Corollary 9. *For $p \in \mathbb{N}$, we have*

$$\sum_{n=1}^{\infty} (n)_{p,\lambda} \left(\frac{1}{1-x} - \sum_{l=0}^n x^l \right) = \frac{x}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right).$$

From (7) and (8) we note that

$$(36) \quad \begin{aligned} \frac{1}{1-x} \text{Li}_{p,\lambda}(x) &= \sum_{l=0}^{\infty} x^l \sum_{k=1}^{\infty} \frac{(-\lambda)^{k-1} (1)_{k,1/\lambda}}{(k-1)! k^p} x^k \\ &= \sum_{n=1}^{\infty} \sum_{k=1}^n \frac{(-\lambda)^{k-1} (1)_{k,1/\lambda}}{k^p (k-1)!} x^n. \end{aligned}$$

For any nonnegative integer r , let D_r denote the linear operator $D_r = \frac{1}{r!} \left(\frac{d}{dx} \right)^r x^r$, so that, for any polynomial $p(x) \in \mathbb{C}[x]$, we have

$$(37) \quad D_r p(x) = \frac{1}{r!} \left(\frac{d}{dx} \right)^r \left[x^r p(x) \right].$$

From (36) and (37) we note that

$$(38) \quad \begin{aligned} D_r F_{n,\lambda}(x) &= \frac{1}{r!} \left(\frac{d}{dx} \right)^r \left[\sum_{k=0}^n S_{2,\lambda}(n, k) k! x^{k+r} \right] \\ &= \frac{1}{r!} \left[\sum_{k=0}^n S_{2,\lambda}(n, k) k! (k+r)_r x^k \right] \\ &= \sum_{k=0}^n S_{2,\lambda}(n, k) \binom{k+r}{k} k! x^k. \end{aligned}$$

Therefore, by (38), we obtain the following theorem.

Theorem 10. *For $n, r \geq 0$, we have*

$$D_r F_{n,\lambda}(x) = \sum_{k=0}^n \binom{k+r}{k} k! S_{2,\lambda}(n, k) x^k.$$

Let $p \in \mathbb{N}$. On the one hand, we have

$$(39) \quad \begin{aligned} \sum_{n=0}^{\infty} \binom{n+r}{r} (n)_{p,\lambda} x^n &= \sum_{n=0}^{\infty} \binom{n+r}{r} x^n \sum_{k=0}^p S_{2,\lambda}(p, k) (n)_k \\ &= \sum_{k=0}^p S_{2,\lambda}(p, k) k! \sum_{n=k}^{\infty} \binom{n+r}{r} \binom{n}{k} x^n \\ &= \sum_{k=0}^p S_{2,\lambda}(p, k) k! \sum_{n=k}^{\infty} \binom{k+r}{k} \binom{n+r}{n-k} x^n \\ &= \sum_{k=0}^p S_{2,\lambda}(p, k) k! \binom{k+r}{r} x^k \sum_{n=0}^{\infty} \binom{n+k+r}{n} x^n \\ &= \sum_{k=0}^p S_{2,\lambda}(p, k) k! \binom{k+r}{r} x^k \left(\frac{1}{1-x} \right)^{r+k+1} \\ &= \left(\frac{1}{1-x} \right)^{r+1} D_r F_{p,\lambda} \left(\frac{x}{1-x} \right). \end{aligned}$$

On the other hand, we also have

$$(40) \quad \left(x \frac{d}{dx} \right)_{p,\lambda} \left(\frac{1}{1-x} \right)^{r+1} = \sum_{n=0}^{\infty} \binom{n+r}{n} \left(x \frac{d}{dx} \right)_{p,\lambda} x^n = \sum_{n=0}^{\infty} \binom{n+r}{n} (n)_{p,\lambda} x^n.$$

Therefore, by (39) and (40), we obtain the following theorem.

Theorem 11. For $p \in \mathbb{N}$, and $r \in \mathbb{Z}$ with $r \geq 0$, we have

$$\left(x \frac{d}{dx}\right)_{p,\lambda} \left(\frac{1}{1-x}\right)^{r+1} = \sum_{n=0}^{\infty} \binom{n+r}{n} (n)_{p,\lambda} x^n = \left(\frac{1}{1-x}\right)^{r+1} D_r F_{p,\lambda} \left(\frac{x}{1-x}\right).$$

By using Theorem 8, we observe that

$$\begin{aligned} (41) \quad \frac{1}{r!} \left(\frac{d}{dx}\right)^r \left[\frac{x^r}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right) \right] &= \frac{1}{r!} \left(\frac{d}{dx}\right)^r \left[\sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) \right] x^{n+r} \\ &= \sum_{n=1}^{\infty} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) \binom{n+r}{r} x^n. \end{aligned}$$

Now, from Theorem 11 and (41) we obtain

$$\begin{aligned} (42) \quad & \sum_{k=1}^{\infty} (k)_{p,\lambda} \left(\left(\frac{1}{1-x}\right)^{r+1} - \sum_{l=0}^k \binom{r+l}{l} x^l \right) \\ &= \sum_{k=1}^{\infty} (k)_{p,\lambda} \sum_{n=k+1}^{\infty} \binom{n+r}{n} x^n = \sum_{n=1}^{\infty} \binom{n+r}{n} x^n \sum_{k=0}^{n-1} (k)_{p,\lambda} \\ &= \sum_{n=1}^{\infty} \binom{n+r}{n} x^n \left(\sum_{k=1}^n (k)_{p,\lambda} - (n)_{p,\lambda} \right) \\ &= \sum_{n=1}^{\infty} \binom{n+r}{r} \left((1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda} \right) x^n - \sum_{n=1}^{\infty} \binom{n+r}{r} (n)_{p,\lambda} x^n \\ &= \frac{1}{r!} \left(\frac{d}{dx}\right)^r \left[\frac{x^r}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right) \right] - \left(x \frac{d}{dx}\right)_{p,\lambda} \left(\frac{1}{1-x}\right)^{r+1}. \end{aligned}$$

Therefore, by (42), we obtain the following differential equation for the degenerate Fubini polynomials.

Theorem 12. For $r \in \mathbb{Z}$ with $r \geq 0$, and $p \in \mathbb{N}$, we have

$$\begin{aligned} & \frac{1}{r!} \left(\frac{d}{dx}\right)^r \left[\frac{x^r}{(1-x)^2} F_{p,\lambda} \left(\frac{x}{1-x}\right) \right] - \left(x \frac{d}{dx}\right)_{p,\lambda} \left(\frac{1}{1-x}\right)^{r+1} \\ &= \sum_{k=1}^{\infty} (k)_{p,\lambda} \left(\frac{1}{(1-x)^{r+1}} - \sum_{j=0}^k \binom{r+j}{j} x^j \right). \end{aligned}$$

3. CONCLUSION

In this paper, we studied the degenerate Fubini and the degenerate Bell polynomials and found some further properties and identities on such polynomials which

had been introduced by Kim-Kim. Especially, we found several expressions for generating functions for the sum $S_{n,\lambda}(p) = (1)_{p,\lambda} + (2)_{p,\lambda} + \cdots + (n)_{p,\lambda}$ of the values of the generalized falling factorials at positive consecutive integers.

As we mentioned earlier, there are various ways of studying special numbers and polynomials, including generating functions, p -adic analysis, umbral calculus, combinatorial methods, differential equations, special functions, probability theory, analytic number theory and operator theory.

It is one of our future projects to continue to explore various degenerate versions of many special polynomials and numbers by making use of such various tools.

REFERENCES

1. ARACI, S. *A new class of Bernoulli polynomials attached to polyexponential functions and related identities*. Adv. Stud. Contemp. Math. (Kyungshang) **31** (2021), no. 2, 195-204.
2. BOYADZHIEV, K. N. *Sums of powers and special polynomials*. Discuss. Math. Gen. Algebra Appl. **40** (2020), no. 2, 275-283.
3. CARLITZ, L. *Degenerate Stirling, Bernoulli and Eulerian numbers*. Utilitas Math. **15** (1979), 51-88.
4. COMTET, L. *Advanced combinatorics. The art of finite and infinite expansions*. Revised and enlarged edition. D. Reidel Publishing Co., Dordrecht, 1974. xi+343 pp. ISBN:
5. FURDUI, O. *Limits, series, and fractional part integrals. Problems in mathematical analysis. Problem Books in Mathematics*. Springer, New York, 2013. xviii+274 pp.
6. JANG, L.C.; KIM, H. *A note on the modified type 2 degenerate poly-Changhee-Genocchi numbers and polynomials*. Adv. Stud. Contemp. Math. (Kyungshang) **31** (2021), no. 3, 325-333.
7. KILAR, N.; SIMSEK, Y. *Identities and relations for Fubini type numbers and polynomials via generating functions and p -adic integral approach*. Publ. Inst. Math. (Beograd) (N.S.) **106** (120) (2019), 113-123.
8. KIM, D. S.; KIM, T. *A note on a new type of degenerate Bernoulli numbers*. Russ. J. Math. Phys. **27** (2020), no. 2, 227-235.
9. KIM, H. K. *Central Lah numbers and central Lah-Bell numbers*. Adv. Stud. Contemp. Math. (Kyungshang) **32** (2022), no. 1, 103-111.
10. KIM, T.; KIM, D. S. *Degenerate Whitney numbers of first and second kind of Dowling lattices*. Russ. J. Math. Phys. **29** (2022), no. 3, 358-377.
11. KIM, T.; KIM, D. S. *On some degenerate differential and degenerate operator*. Russ. J. Math. Phys. **29**(2022), no. 1, 37-47.
12. KIM, T.; KIM, D. S. *Some identities on truncated polynomials associated with degenerate Bell polynomials*. Russ. J. Math. Phys. **28** (2021), no. 3, 342-355.
13. KIM, T.; KIM, D. S. *Note on the degenerate gamma function*, Russ. J. Math. Phys. **27** (2020), no. 3, 352-358.
14. KIM, T.; KIM, D. S.; DOLGY, D. V. *On partially degenerate Bell numbers and polynomials*. Proc. Jangjeon Math. Soc. **20** (2017), no. 3, 337-345.

15. KIM, T.; KIM, D. S.; DOLGY, D. V.; PARK, J.-W. *Degenerate binomial and Poisson random variables associated with degenerate Lah-Bell polynomials*. Open Math. **19** (2021), no. 1, 1588-1597.
16. KIM, T.; KIM, D. S.; LEE, H.; KWON, J. *Degenerate binomial coefficients and degenerate hypergeometric functions*. Adv. Difference Equ. 2020, (2020:115).
17. KIM, T.; KIM, D. S.; KIM, H. Y.; KWON, J. *Some results on degenerate Daehee and Bernoulli numbers and polynomials*. Adv. Difference Equ. 2020, (2020:311).
18. ROMAN, S. The umbral calculus. Pure and Applied Mathematics, 111. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984. x+193 pp. ISBN: 0-12-594380-6
19. SHARMA, S. K.; KHAN, W. A.; ARACI, S.; AHMED, S. S. *New construction of type 2 degenerate central Fubini polynomials with their certain properties*. Adv. Difference Equ. 2020, Paper No. 587, 11 pp.
20. SIMSEK, Y. *Construction of generalized Leibnitz type numbers and their properties*. Adv. Stud. Contemp. Math. (Kyungshang) **31** (2021), no. 3, 311-323.

Taekyun Kim

Department of Mathematics,
Kwangwoon University,
Seoul 139-701, Republic of Korea,
E-mail: tkkim@kw.ac.kr

(Received 10. 03. 2020.)

(Revised 02. 10. 2022.)

Dae San Kim

Department of Mathematics,
Sogang University,
Seoul 121-742, Republic of Korea,
E-mail: dskim@sogang.ac.kr