



HAL
open science

Albanese map of special manifolds: a correction

Frederic Bruno Campana

► **To cite this version:**

| Frederic Bruno Campana. Albanese map of special manifolds: a correction. 2021. hal-03344242

HAL Id: hal-03344242

<https://hal.science/hal-03344242>

Preprint submitted on 14 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ALBANESE MAP OF SPECIAL MANIFOLDS: A CORRECTION

FRÉDÉRIC CAMPANA

1. ABSTRACT

We show that any fibration of a ‘special’ compact Kähler manifold X onto an Abelian variety has no multiple fibre in codimension one. This statement strengthens and extends previous results of Kawamata and Viehweg when $\kappa(X) = 0$. This also corrects the proof given in [2], 5.3 which was incomplete.

2. INTRODUCTION

The following statement is given in [2], Proposition 5.3:

Theorem 2.1. *Let $a_X : X \rightarrow A_X$ is the Albanese map of X , assumed to be special. Then a_X is onto, has connected fibres, and no multiple fibre in codimension one.*

Recall ([2], definition 2.1 and Theorem 2.27) that X is special if $\kappa(X, L) < p$ for any $p > 0$, and any rank-one coherent subsheaf of Ω_X^p . This implies that X has no surjective meromorphic $g : X \dashrightarrow Z$ map onto a manifold Z of general type and positive dimension p , and more generally exactly means that the ‘orbifold base’ (Z, D_g) of such a g , constructed out of its multiple fibres, is never of ‘general type’. See §.2 below for some more details.

While the proofs given there for the first two properties (which generalise earlier results by Y. Kawamata and Kawamata-Viehweg in the case when $\kappa(X) = 0$) are complete, the proof of the third property, even in the projective case, is not (as pointed out to me by K. Yamanoi and E. Rousseau¹). The aim of this note is to correct it in the projective case (but only partially in the compact Kähler case) by using the main result of [3], itself based on [1] (or, alternatively, [4]).

Theorem 2.2. *Let $f : X \rightarrow A$ be a holomorphic map from the connected, compact, Kähler manifold to a compact complex torus A . If X is special, then:*

Date: September 14, 2021.

¹The problem comes from the potential multiplicity-one exceptional divisors of a_X which are no longer f -exceptional if $g = q \circ f$, where $q : A \rightarrow B$ is a non-trivial torus quotient sending D_f to an ample divisor of B . We overcome this difficulty in the second step of the proof, by cutting the fibration by means of Poincaré reducibility. The first step is the same as in [2].

1. f is surjective,
2. If f is the Albanese map of X , then f has connected fibres, and:
3. f has no multiple fibre² in codimension one if the fibres of f are connected, and if A is an Abelian variety (equivalently: $D_f = 0$ if (A, D_f) is the ‘orbifold base’ of f).

Remark 2.3. 1. Although the assumption that A is an Abelian variety in Claim 3 is certainly not necessary, our proof uses [1] (or alternatively [4]), presently known only for A projective. Even with a Kähler version of [1] or [4], we could not treat the Kähler case in general with the present arguments, because we use Poincaré reducibility.

2. An easy case of Claim 3, not covered by Theorem 2.2 is when the algebraic dimension $a(X)$ of X vanishes (and more generally, if $a(A_X) = 0$, A_X being the Albanese torus of X), because then A_X does not contain any effective divisor. The argument of the second step of the proof below shows that the conclusion of Theorem 2.2.(3) still holds true if $A = T \times V$, where T is a compact torus of algebraic dimension zero, and V an abelian variety.

3. Claims 1 and 2, and their proofs, still apply when X is ‘weakly special’, which means that no finite étale cover X' of X has a surjective meromorphic fibration onto a positive-dimensional projective manifold of general type. Weak-specialness coincides with specialness for curves and surfaces, but differ from dimension 3 on, by examples of Bogomolov-Tschinkel.

3. MULTIPLE FIBRES OF MAPS TO COMPLEX TORI.

If $f : X \rightarrow Y$ is a fibration, that is: a holomorphic map with connected fibres onto a (smooth) compact complex manifold Y , we define its ‘orbifold base’ D_f as follows: for every prime divisor $E \subset Y$, let $f^*(E) := \sum_{k \in K} t_k \cdot D_k + R$, where the D'_k s are the pairwise distinct prime divisors of X surjectively mapped by f onto E , while R is an effective f -exceptional divisor of X . The multiplicity³ $m_f(E)$ of the generic fibre of f over E is then defined as: $m_f(E) := \inf\{t_k, k \in K\}$, and then $D_f := \sum_{E \subset Y} (1 - \frac{1}{m_f(E)}) \cdot E$, which is an effective \mathbb{Q} -divisor of Y (note indeed that this sum is finite since $m_f(E) = 1$ for all, but finitely many of the E 's).

Proof. (of Theorem 2.2)

Claim 1. Assume by contradiction that $f(X) := Z \neq A$. After Ueno’s theorem ([7], p. 120), there is a quotient torus $q : A \rightarrow B$ such that $q(Z) \subset B$ is of dimension $p > 0$, and of general type. The

²We use here the ‘inf’ multiplicities, not the more classical ‘gcd’ version, which would give a weaker result.

³We use here the ‘inf’ multiplicity instead of the usual ‘gcd’ multiplicity for reasons explained in [2].

composed map $g := q \circ f : X \rightarrow Z$ thus contradicts the specialness of X , since $\kappa(X, g^*(K_Z)) = p$ and $g^*(K_Z) \subset \Omega_X^p$.

Claim 2. Let Z be a smooth model of the Stein factorisation Z_0 of f . We replace X by Z , which is still special since dominated by X . We may thus assume that f is generically finite. By [6], Theorem 23, replacing Z by a suitable finite étale cover which is still special by [2], §.5.5, we may assume that Z fibres over over a manifold of general type if Z is not birational to A . This again contradicts the specialness of Z and thus of X .

Claim 3. Assume that $D_f \neq 0$. We shall first treat the case when D_f is ample, and reduce to this case in a second step.

- Let thus D_f be ample on A . By a flattening of f , followed by suitable blow-ups of X and A , we may assume (see [2], Lemma 1.3) that $f = v \circ f'$, where $v : A' \rightarrow A$ is bimeromorphic with A' smooth, and $f' : X \rightarrow A'$ is a fibration such that its orbifold base $D' := D_{f'} = \overline{D} + E'$, with E' effective and v -exceptional, \overline{D} is the strict transform of D_f in A' . This shows that in particular $f'^*(K_{A'} + \overline{D}) \subset \Omega_{X'}^p, p := \dim(A)$. The following lemma 3.1 shows that the line bundle $K_{A'} + \overline{D}$ has Kodaira dimension $\dim(A) = p$, since D_f is ample on A . This contradicts the specialness of X if D_f is assumed to be ample.

Lemma 3.1. ([2], 1.14) *Let $v : Y' \rightarrow Y$ be a bimeromorphic map between compact connected complex manifolds. Let D be an effective \mathbb{Q} -divisor on Y , and let \overline{D} be its strict transform on Y' . Then:*

1. $K_{Y'} + \overline{D} = v^*(K_Y + \varepsilon.D) + E', \forall \varepsilon > 0$ small enough, E' effective.
2. If $\kappa(Y) \geq 0$, then $\kappa(Y', K_{Y'} + \overline{D}) = \kappa(Y, K_Y + D)$.

Proof. Claim 1. Let the rational numbers a_i, b_i be defined as follows: $v^*(D) = \overline{D} + \sum_i b_i.E_i, K_{Y'} = v^*(K_Y) + \sum_i a_i.E_i$, where the E_i 's are the exceptional divisors of v . Then: $a_i > 0, b_i \geq 0, \forall i$. Then:

$K_{Y'} + \overline{D} = v^*(K_Y) + \sum a_i.E_i + (1 - \varepsilon).\overline{D} + v^*(\varepsilon.D) - (\varepsilon.\sum b_i.E_i) = v^*(K_Y + \varepsilon.D) + (1 - \varepsilon).\overline{D} + \sum (a_i - \varepsilon.b_i).E_i$. We thus get the first claim when $0 < \varepsilon \leq \min\{1, \frac{a_i}{b_i}, \forall i\}$, with $E' := (1 - \varepsilon).\overline{D} + \sum (a_i - \varepsilon.b_i).E_i$.

Claim 2. Let $\varepsilon > 0$ be as above. Since K_Y is assumed to be \mathbb{Q} -effective, so is $(1 - \varepsilon).v^*(K_Y)$, and so $K_{Y'} + \overline{D} = \varepsilon.v^*(K_Y + D) + E''$, with: $E' := E' + (1 - \varepsilon).v^*(K_Y)$, which is \mathbb{Q} -effective. \square

Remark 3.2. *The main property used in Lemma 3.1 is that $a_i > 0, \forall i$, i.e: the lift of the canonical sheaf under a modification of a smooth manifold vanishes on the exceptional divisor. This property does not hold true for sheaves of forms of degree less than the dimension. We thus need to use a less direct route, which requires deeper ingredients.*

We shall now reduce to the case when D_f is ample.

- Let $L := \mathcal{O}_A(m.D_f)$ be the line bundle on A with a section vanishing on some integral multiple $m.D_f$ of the orbifold base of f . Let

$T \subset \text{Aut}^0(A)$ be the connected component of the group of translations of A preserving L . Let $q : A \rightarrow B := A/T$ be the quotient map. Then $D_f = \frac{1}{m} \cdot q^*(D)$ for some *ample* effective divisor D on B (see, for example [5], Théorème 5.1). Since $D_f \neq 0$, $p := \dim(B) > 0$.

We use Poincaré reducibility to reduce to the case that $A = B \times B'$, where $B' \rightarrow T$ is a suitable finite étale cover of T . This indeed amounts to replace X and D_f by the corresponding finite étale covers, preserving all of our hypothesis, in particular the specialness of X (by [2], §.5.5, again).

We now consider the composed fibrations $g := q \circ f : X \rightarrow B$, and $h := q' \circ f : X \rightarrow B'$, where $q' : A \rightarrow B'$ is the second projection. By [3], 2.4, the general fibre $X_{b'} := h^{-1}(b')$, $b' \in B'$ of g' , is special⁴. Let $f_{b'} : X_{b'} \rightarrow A_{b'} := B \times \{b'\}$ be the restriction of f to $X_{b'}$. The orbifold base $(A_{b'}, D_{f_{b'}})$ of $f_{b'}$ is then nothing, but $(A, D_f) \cap A_{b'}$, and is thus ample on $A_{b'} \cong B$. By the first part of the proof, this contradicts the specialness of $X_{b'}$ if $D_{f_{b'}} \neq 0$. Thus $D_f \cap A_{b'} = 0$, and so $D_f = 0$. \square

REFERENCES

- [1] C. Birkar-J. Chen. Varieties fibered over Abelian varieties with fibres of Log-general type. *Adv. Math.* 270 (2015), 206-222.
- [2] F. Campana. Orbifolds, special varieties and classification theory. *Ann. Inst. Fourier*, tome 54 (2004), 499-630.
- [3] F. Campana-B. Claudon. Quelques propriétés de stabilité des variétés spéciales. *Math. Zeitschrift* 283 (2016), 581-599.
- [4] J. Cao-M. Păun. Kodaira dimension of algebraic fibre spaces over abelian varieties. *Inv. Math.* 207 (2017), 345-387.
- [5] O. Debarre. *Tores et variétés Abéliennes complexes. Cours spécialisés 6.* SMF et EDP Sciences 1999.
- [6] Y. Kawamata. Characterisation of Abelian varieties. *Comp. Math.* 43 (1981), 253-276.
- [7] K. Ueno. *Classification theory of algebraic varieties and compact complex analytic spaces.* Lect.Notes.Math. 439. Springer Verlag (1975).

UNIVERSITÉ LORRAINE, NANCY,
Email address: frederic.campana@univ-lorraine.fr

⁴The statement of [3] is only given for X projective, and f the Albanese map of X , but it is easy to check that the proof, derived from [1], or [4], through [3], 2.6, applies for X compact and any map to an Abelian variety.