



available at www.sciencedirect.com



journal homepage: www.elsevier.com/locate/jjcc



Corrigendum

Corrigendum to ‘Investigation of optimal anticoagulation strategy for stroke prevention in Japanese patients with atrial fibrillation—The J-RHYTHM Registry study design’ [J. Cardiol. 57 (2011) 95–99]

Hirotsugu Atarashi (MD, PhD, FJCC)^{a,*}, Hiroshi Inoue (MD, PhD, FJCC)^b,
Ken Okumura (MD, PhD, FJCC)^c, Takeshi Yamashita (MD, PhD, FJCC)^d,
Hideki Origasa (PhD)^e, for the J-RHYTHM Registry Investigators

^a Department of Internal Medicine, Nippon Medical School Tama-Nagayama Hospital, Tokyo, Japan

^b The Second Department of Internal Medicine, University of Toyama, Graduate School of Medicine, Toyama, Japan

^c Cardiology, Respiratory Medicine and Nephrology, Hirosaki University Graduate School of Medicine, Aomori, Japan

^d The Cardiovascular Institute, Tokyo, Japan

^e Division of Biostatistics and Clinical Epidemiology, University of Toyama, Graduate School of Medicine, Toyama, Japan

The author regrets that in the original printing of the above article on page 96 some information was printed incorrectly. The correct information is as follows:

- 1) In the paragraph of ‘‘Study Population’’ in the line 4, from ‘‘who had maintained a sinus rhythm for more than 1 year’’ to ‘‘, but those patients who have maintained a sinus rhythm for more than 1 year, as well as in-patient were excluded from the study’’.
- 2) In the paragraph of ‘‘Study design and patient enrollment’’, in the last sentence and last line, from ‘‘January 2010 and July 2010’’ to ‘‘January 2009 and July 2009’’.

DOI of original article: [10.1016/j.jjcc.2010.09.002](https://doi.org/10.1016/j.jjcc.2010.09.002).

* Corresponding author at: Nippon Medical School Tama-Nagayama Hospital, Division of Cardiology, 1-7-1 Nagayama, Tama-city, Tokyo 206-8512, Japan. Tel.: +81 42 371 2111; fax: +81 42 371 2100.

E-mail address: atarashi@nms.ac.jp (H. Atarashi).